

Positioning a catheter with an expandable treatment balloon in the prostatic urethra

Block 100

Massaging substantially the entire circumference of the lumen in the upper portion of the prostatic urethra extending between the bladder neck and above the verumontanum

Block 115

Internally massaging at least a portion of the prostatic urethra by repetitively outwardly expanding and then contracting the treatment balloon

Block 110

Concurrently heating at least a portion of the prostatic urethra during the internal massage

Block 120





Figure 4

0.1 ocm ("L1") - up to 6.0 cm ("L2")







Figure 5B

Time






FIGURE 7

Average:	10.30	4.35	7.40	7.23	3.23	5.00	5.43	3.14	3.8
20	16	10	8			 			
19	10	2	7	1	 	-			
18	13	0	- 10	14	0	10			<u> </u>
17	3	0	1	 		 	- 		
16	15	4	11	+	 				
15	18	7.	9					-	
14	14	0	9	1	<u> </u>				
13	15	10	8	15	10	8			
12	10	6	10	8	2	5	 	-	6
11	9	4	7	11	4	7	10	4	
10	9	2	7	- -	 	 		 	
9	3	4	6	2	0	1	2	1	<u> </u>
8	5	2	3	1	2	2		 	
7	8	- 6	8	4	5 .	3		 	-
6	7	5	10	7	1 0	4		 	
5	9	5	7	6	. 2	5	6	2	-
4	4	2	2	4	3	3	2	2	-
3	11	1	5	5	+ 1	3	10	1	-
2	16	10	12	15	9	11	16	8	<u> </u>
1	11	7	8	2	4	3	2	4	
	· P	US	QL	Р	T us	T QL	P	CPSI	, ,
of Patients	Baseline CPSI			Comments - 1 month			Comments - 3 mor		
- 6 0	i	_		1 1			1 1		


P = Pain

US = Urinary System

QL = Quality of Life


Figure 8

WIT and Chronic Nonbacterial Prostatitis CLINICAL PERFORMANCE



THE POWER OF PRACTICAL TECHNOLOGIES

WIT and Chronic Nonbacterial Prostatitis CLINICAL PERFORMANCE



WIT and Chronic Nonbacterial Prostatitis CLINICAL PERFORMANCE

